

Table 4 Comparison of practices of lecture and videotape groups to reduce exposure of their children to pesticides at home, pre-training, and 2 weeks and 1 month after the training

Practice	Pre-training		2 weeks after		1 month after	
	Lecture	Video	Lecture	Video	Lecture	Video
	(n = 165) %	(n = 132) %	(n = 165) %	(n = 132) %	(n = 165) %	(n = 132) %
Apply pesticides as needed only	41.0	43.0	81.0	88.0	79.0	92.0
Avoid use of pesticides on children's bodies	83.0	82.0	99.0	97.0	99.0	97.0
Never allow children to use pesticides	61.0	63.0	92.0	95.0	91.0	96.0
Keep children's clothes and toys away during pesticide application	43.0	40.0	92.0	94.0	90.0	93.0
Keep food and drinks away during pesticide application	76.0	78.0	94.0	94.0	93.0	95.0
Teach children to wash hands before eating and after playing	58.0	56.0	87.0	92.0	85.0	90.0
Wash vegetables and fruits before eating	89.0	88.0	99.0	99.0	100.0	100.0
Never use empty pesticide containers in other purposes	69.0	71.0	89.0	96.0	85.0	95.0
Keep pesticides away from children	54.0	64.0	90.0	92.0	88.0	89.0

n = total number of participants.