

**Table 2 Association of health and sexual hygiene practices with urinary tract infection for cases ( $n = 100$ ) and matched controls ( $n = 150$ )**

Practice	Cases				Controls				OR (95% CI)
	Yes		No		Yes		No		
	No.	%	No.	%	No.	%	No.	%	
Not washing genitals precoitus	47	47.0	53	53.0	98	65.8	51	34.2	2.16 (1.29–3.63)
Husband not washing genitals precoitus	52	52.0	48	48.0	110	73.3	40	26.7	2.53 (1.48–4.32)
Not washing genitals postcoitus	84	84.0	16	16.0	143	95.3	7	4.7	2.89 (1.53–9.80)
Washing genitals from back to front	60	60.0	40	40.0	120	81.6	27	18.4	2.96 (1.66–5.28)
Not voiding urine postcoitus	51	51.0	49	49.0	135	90.0	15	10.0	8.62 (6.66–16.66)
Delay in voiding urine (voluntary)	69	69.7	30	30.3	52	34.7	98	65.8	4.33 (2.51–7.47)
Not drying after voiding	39	39.0	61	61.0	104	69.3	46	30.7	3.53 (1.61–6.21)
Not drinking plenty of liquids	32	32.0	68	68.0	78	54.2	66	45.8	2.51 (1.45–4.24)

*Data missing for some variables.*

*OR = odds ratio; CI =confidence interval.*