

Table 3 Results of susceptibility testing of *Pseudomonas aeruginosa* (n = 45 isolates), and minimum inhibitory concentration (MIC) of each antibiotic used

Antibiotic	Sensitive			Intermediate			Resistant		
	No.	%	MIC ( $\mu$ g/mL)	No.	%	MIC ( $\mu$ g/mL)	No.	%	MIC ( $\mu$ g/mL)
Amikacin	45	100.0	$\leq 16$	0	–	32	0	–	$\geq 64$
Aztreonam	36	80.0	$\leq 8$	8	17.8	16	1	2.2	$\geq 32$
Cefepime	45	100.0	$\leq 8$	0	–	16	0	–	$\geq 32$
Ceftazidime	45	100.0	$\leq 8$	0	–	16	0	–	$\geq 32$
Ciprofloxacin	45	100.0	$\leq 1$	0	–	2	0	–	$\geq 4$
Gentamicin	45	100.0	$\leq 4$	0	–	8	0	–	$\geq 16$
Imipenem	44	97.8	$\leq 4$	0	–	8	1	2.2	$\geq 16$
Meropenem	44	97.8	$\leq 4$	0	–	8	1	2.2	$\geq 16$
Piperacillin	45	100.0	$\leq 64$	0	–	–	0	–	$\geq 128$
Ticarcillin/clavulanic acid	36	80.0	$\leq 64/< 2$	0	–	–	9	20.0	$\geq 128/\geq 2$
Tobramycin	45	100.0	$\leq 4$	0	–	8	0	–	$\geq 16$