

Table 3 Prevalence of other risk factors among patients with hypertension and the controls

Risk factor	Cases		Controls		P-value	OR (95% CI)
	No.	%	No.	%		
<i>Smoking</i>						
Non-smokers	163	73.8	183	82.8	< 0.01	
Ex-smokers	25	11.3	8	3.6		
Smokers	33	14.9	30	13.6		1.23 ^a (1.06-1.59)
<i>Body mass index (kg/m²)</i>						
< 25	43	19.5	74	33.5	< 0.01	
25-29.9	96	43.4	80	36.2		
≥ 30	82	37.1	67	30.3		2.10 ^b (1.28-3.46)
<i>Physical activity level</i>						
Active	21	9.5	36	16.3	< 0.01	
Irregular	24	10.9	41	18.6		
Inactive	176	79.6	144	65.2		2.10 ^c (1.28-3.74)
<i>Diabetes mellitus</i>						
Yes	58	26.2	33	14.9	< 0.01	
No	163	73.8	188	85.1		2.03 (1.26-3.27)
<i>Family history of hypertension</i>						
Yes	142	64.3	108	48.9	< 0.01	
No	79	35.7	113	51.1		1.88 (1.29-2.75)
Total	221	100.0	221	100.0		

^aSmokers versus non-smokers.^bBody mass index ≥ 30 kg/m² versus < 25 kg/m².^cInactive versus active.

OR = odds ratio; CI = confidence interval.