

Table 2 Antibiotic susceptibility rates of nosocomial bacterial isolates (n = 556)

Antibiotic	<i>Escherichia coli</i> (n = 238)		<i>Acinetobacter</i> spp. (n = 95)		<i>Klebsiella</i> spp. (n = 92)		<i>Pseudomonas aeruginosa</i> (n = 79)		<i>Enterobacter</i> spp. (n = 42)		Others (n = 10)		Total (n = 556)	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Gentamicin	125	52.5	14	14.7	48	52.2	34	43.0	26	61.9	8	80.0	285	51.3
Ciprofloxacin	90	37.8	27	28.4	48	52.2	43	54.4	31	73.8	7	70.0	264	47.5
Amikacin	81	34.0	15	15.8	29	31.5	24	30.4	25	59.5	4	40.0	178	32.0
Ceftazidime	75	31.5	5	5.3	32	34.8	24	30.4	22	52.4	7	70.0	165	29.7
Ceftriaxone	64	26.9	2	2.1	30	32.6	6	7.6	15	35.7	5	50.0	123	22.1
Nitrofurantoin	70	29.4	2	2.1	4	4.3	0	0.0	1	2.4	0	0.0	77	13.8
Nalidixic acid	34	14.3	0	0.0	14	15.2	1	1.3	0	0.0	1	10.0	50	9.0
Cefazolin	21	8.8	2	2.1	13	14.1	5	6.3	1	2.4	3	30.0	45	8.1
Tobramycin	8	3.4	6	6.3	4	4.3	12	15.2	12	28.6	1	10.0	43	7.7
Piperacillin/tazobactam	5	2.1	4	4.2	0	0.0	9	11.4	6	14.3	0	0.0	24	4.3
Piperacillin	1	0.4	2	2.1	0	0.0	6	7.6	5	11.9	0	0.0	14	2.5
Cefuroxime	2	0.8	0	0.0	2	2.2	2	2.5	0	0.0	1	10.0	7	1.3