

**Table 5 Role of health-care providers in advising smokers to use smoking cessation clinics or hotline services in relation to sociodemographic characteristics**

Variable	Referred to cessation clinic or hotline by health-care provider					
	No		Yes		Have not visited health-care provider recently	
	No.	%	No.	%	No.	%
<b>Total</b>	680	65.9	205	19.9	147	14.2
<b>Sex</b>						
Male	600	66.4	176	19.5	128	14.2
Female	80	62.5	29	22.7	19	14.8
			$\chi^2 = 0.865; P = 0.649$			
<b>Age (years)</b>						
18–29	207	69.6	45	15.2	44	14.9
30–39	229	68.2	61	18.2	46	13.7
40–49	150	59.8	60	23.9	41	16.3
50–59	56	67.5	19	22.9	8	9.6
60+	39	57.4	21	30.9	8	11.8
			$\chi^2 = 15.97; P = 0.043$			
<b>Education</b>						
Elementary or less	111	66.9	19	11.4	36	21.7
Preparatory to high school	388	64.2	124	20.5	92	15.2
Diploma and above	181	69.3	61	23.4	19	7.3
			$\chi^2 = 24.19; P < 0.001$			
<b>Monthly income (JD)<sup>a</sup></b>						
< 300	339	67.4	84	16.7	80	15.9
300–599	262	67.2	77	19.7	51	13.1
600+	78	57.4	43	31.6	15	11.0
			$\chi^2 = 16.13; P = 0.003$			

<sup>a</sup>6 respondents refused to declare their income.