

Table 2 Resistance pattern of *Acinetobacter baumannii* isolates from different clinical specimens

Antibiotic	Swab (n = 57)		Blood (n = 24)		Sputum (n = 28)		Urine (n = 7)		Total (n = 116)		P-value
	No.	%	No.	%	No.	%	No.	%	No.	%	
Ceftazidime	56	98.2	23	95.8	28	100.0	7	100.0	114	98.3	0.692
Ciprofloxacin	57	100.0	18	75.0	28	100.0	7	100.0	110	94.8	0.000
Ceftriaxone	57	100.0	24	100.0	28	100.0	7	100.0	116	100.0	-
Colistin	1	1.8	1	4.2	0	0.0	0	0.0	2	1.7	0.692
Cefotaxime	57	100.0	24	100.0	28	100.0	7	100.0	116	100.0	-
Cefepime	56	98.2	23	95.8	28	100.0	7	100.0	114	98.3	0.692
Gentamicin	53	93.0	17	70.8	26	92.9	6	85.7	102	87.9	0.034
Imipenem	56	98.2	22	91.7	28	100.0	7	100.0	113	97.4	0.237
Levofloxacin	57	100.0	18	75.0	28	100.0	7	100.0	110	94.8	0.000
Minocycline	12	21.1	3	12.5	12	42.9	4	57.2	31	26.7	0.015
Piperacillin	56	98.2	23	95.8	28	100.0	7	100.0	114	98.3	0.692
Ampicillin/sulbactam	51	89.5	24	100.0	22	78.6	7	100.0	104	89.7	0.063
Trimethoprim/sulfamethoxazole	47	82.5	20	83.3	18	64.3	3	42.9	88	75.9	0.040
Tetracycline	42	73.7	14	58.3	26	92.9	7	100.0	89	76.7	0.011
Ticarcillin	57	100.0	24	100.0	28	100.0	7	100.0	116	100.0	-
Tobramycin	42	73.7	16	66.7	25	89.3	6	85.7	89	76.7	0.217
Piperacillin/tazobactam	55	96.5	22	91.7	28	100.0	7	100.0	112	96.6	0.396