

Table 2 Demographic and clinical characteristics of cutaneous leishmaniasis patients with *Leishmania tropica* and their responses to Glucantime[®]

Variable	Cure		Failure		P-value ^a
	No.	%	No.	%	
Sex					
Male	74	97.4	2	2.6	0.662
Female	62	95.4	3	4.6	
Age group (years)					
< 10	28	87.5	4	12.5	0.01
≥ 10	108	99.1	1	0.9	
Type of injection					
Intramuscular	46	90.2	5	9.8	0.005
Intralesional	90	100.0	0	0.0	
Intramuscular injection by age group (years)					
< 10	13	77.8	4	22.2	0.037
≥ 10	33	97.0	1	3.0	
Duration of intramuscular- therapy (days)					
< 10	1	100.0	0	0.0	0.416
10–14	3	75.0	1	25.0	
≥ 15	42	91.3	4	8.7	
No. of intralesional injections					
1–4	5	100.0	0	0.0	1.0
5–8	18	100.0	0	0.0	
9–12	118	100.0	0	0.0	
No. of lesions					
≤ 3	127	97.7	3	2.3	0.049
> 3	9	81.8	2	18.2	
Location of lesions					
Face and neck	42	93.3	3	6.7	0.465
Extremities	88	97.8	2	2.2	
Both	6	100.0	0	0.0	
Residency					
Urban	89	95.7	4	4.3	0.661
Rural	47	97.9	1	2.1	

^aFisher exact test.