

Table 1 Demographic characteristics and influenza prevalence among patients with ILI and SARI, Morocco, 2014–2015

Characteristic	ILI (<i>n</i> = 273) No. (%)	SARI (<i>n</i> = 167) No. (%)	Total (<i>n</i> = 440) No. (%)
Age			
0–23 months	5 (1.8)	71 (42.5)	76 (17.3)
2–5 years	19 (7.0)	13 (7.8)	32 (7.3)
6–15 years	50 (18.3)	14 (8.4)	64 (14.6)
16–49 years	112 (41.0)	28 (16.8)	140 (31.8)
50–64 years	59 (21.6)	17 (10.2)	76 (17.3)
≥ 65 years	16 (5.9)	15 (9.0)	31 (7.1)
Unknown	12 (4.4)	9 (5.4)	21 (4.8)
Sex			
Female	155 (56.7)	81 (48.5)	236 (53.6)
Influenza subtype			
A (H1N1)pdm09	24 (8.8)	14 (8.4)	38 (8.6)
A (H3N2)	22 (8.1)	3 (1.8)	25 (5.7)
A unsubtypeable	NA	3 (1.8)	3 (0.7)
B	113 (41.4)	22 (13.2)	135 (30.7)
Negative	114 (41.8)	125 (74.9)	239 (54.3)

ILI – influenza-like illness; SARI – severe acute respiratory illness.