

Table 4 Perceived need for education about sexually transmitted infections (STIs) among the midwives

STI	Very low need	Low need	Moderate need	High need	Very high need	Mean score (SD)
	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	
HIV/AIDS	27 (18.8)	35 (24.3)	46 (31.9)	26 (18.1)	10 (6.9)	2.70 (1.17)
Genital herpes	10 (6.9)	17 (11.8)	63 (43.8)	41 (28.5)	13 (9.0)	3.21 (1.0)
Genital warts	10 (6.9)	18 (12.5)	66 (45.8)	38 (26.4)	12 (8.3)	3.17 (0.99)
Hepatitis B	15 (10.4)	35 (24.3)	52 (36.1)	32 (22.2)	10 (6.9)	2.91 (1.08)
Gonorrhoea	5 (3.5)	23 (16.0)	50 (34.7)	52 (36.1)	14 (9.7)	3.33 (0.97)
Chlamydia	11 (7.6)	26 (18.1)	56 (38.9)	41 (28.5)	10 (6.9)	3.09 (1.02)
Syphilis	4 (2.8)	18 (12.5)	60 (41.7)	50 (34.7)	12 (8.3)	3.33 (0.90)
Chancroid	7 (4.9)	13 (9.0)	45 (31.3)	62 (43.1)	17 (11.8)	3.48 (0.98)
Granuloma inguinale	6 (4.2)	15 (10.4)	44 (30.6)	57 (39.6)	22 (15.3)	3.51 (1.01)
Trichomoniasis	27 (18.8)	32 (22.2)	43 (29.9)	30 (20.8)	12 (8.3)	2.78 (1.21)
Candidiasis	30 (20.8)	31 (21.5)	43 (29.9)	28 (19.4)	12 (8.3)	2.73 (1.23)
Total	6 (4.2)	28 (19.4)	65 (45.1)	36 (25.0)	9 (6.3)	34.23 (8.36)

SD = standard deviation.