

**Table 1 Coagulation parameters in patients with severe COVID-19.**

Variable	All patients (n=74) Mean±SD	Survivors (n=35) Mean±SD	Non-survivors (n=39) Mean±SD, P	Patients without comorbidities (n=23) Mean±SD	Patients with comorbidities (n=51) Mean±SD, P
<b>Age (years)</b>	65.1±17.1 (23-90)	63.6±18.7	66.5±15.5, 0.48	55.2±16	69.4±15.8, 0.001
<b>Sex (male/female)</b>	44/30	21/14 (47.7/46.7%)	23/16 (52.3/53.3%)	17/6 (38.6/20%)	27/24 (61.4/80%)
PLT counts (×10 <sup>3</sup> /μL)	207.8±111.5 (13.5-707)	223.9±124.1	193±9.8, 0.24	218.7±117.7	186.9±97.3, 0.22
PT (sec)	15.6±3.9 (12.5-35.4)	15.1±2.8	16±4.6, 0.29	14.4±1.6	16.1±4.4, 0.024
aPTT (sec)	39.1±9.2 (25-68)	39±10.5	39.1±8.2, 0.98	38.2±11.2	39.4±8.5, 0.68
INR	1.42±0.62 (1-5.1)	1.4±0.47	1.5±0.72, 0.39	1.22±0.22	1.50±0.7, 0.02
D-dimer (μg/ml)	5.1±7.3 (0.1-25) n=16	4.8±9.9	5.3±5.9, 0.92	5.1±7	5.1±7.6, 0.99
Fibrinogen (g/L)	3.37±1.80 (135-751) n=16	3.38±2.39	3.36±1.28, 0.98	2.69	3.45±1.9, 0.71

PLT=platelet; PT=prothrombin time; aPTT=activated partial thromboplastin time; INR=international normalized ratio