

Table 5 DALYs attributable to inadequate physical activity by age, sex and cancer site, Tunisia, 2019

Age, in years	Post-menopausal breast			Lung			Colon/rectum			Endometrium			Stomach			Kidney			Bladder			Oesophagus		
	AD	PAF	OD	AD	PAF	OD	AD	PAF	OD	AD	PAF	OD	AD	PAF	OD	AD	PAF	OD	AD	PAF	OD	AD	PAF	OD
Men																								
35-44	NA	NA	NA	444	16.7	2 663	98	11.6	851	NA	NA	NA	103	16.7	618	18	11.1	160	36	17.6	206	24	19.5	120
45-64	NA	NA	NA	5 022	17.0	29 546	656	11.9	5 536	NA	NA	NA	662	17.0	3 895	135	11.3	1200	491	18.0	2728	210	19.9	1059
≥ 65	NA	NA	NA	3 946	17.3	22 871	745	12.0	6 183	NA	NA	NA	722	17.3	4 165	128	11.5	1112	984	18.3	5369	64	19.8	324
Total	NA	NA	NA	9 412	17.1	55 080	1499	11.9	12 570	NA	NA	NA	1487	17.1	8 678	281	11.4	2472	1511	18.2	8303	298	19.8	1503
Women																								
35-44	NA	NA	NA	93	16.3	566	152	11.9	1 280	23	19.2	120	119	17.3	684	11	11.5	94	10	18.3	55	8	20.2	38
45-64	1690	12.8	13 237	482	16.6	2 894	727	12.3	5 886	166	19.6	847	373	17.6	2 119	52	11.7	442	75	18.6	405	55	20.6	269
≥ 65	902	12.9	6 988	439	16.9	2 601	736	12.5	5 865	161	19.9	811	411	17.9	2 294	51	11.9	424	152	18.9	806	63	20.9	300
Total	2592	12.8	20 225	1 014	16.7	6 061	1615	12.4	13 032	350	19.7	1778	903	17.7	5 097	114	11.8	960	237	18.8	1266	126	20.7	607
Both																								
35-44	NA	NA	NA	537	16.6	3 229	250	11.7	2 132	23	19.2	120	222	17.0	1 302	29	11.2	254	46	17.7	261	32	19.7	158
45-64	1690	12.8	13 237	5 504	17.0	32 440	1383	12.1	11 422	166	19.6	847	1035	17.2	6 014	187	11.4	1642	566	18.1	3133	265	20.0	1328
≥ 65	902	12.9	6 988	4 385	17.2	25 472	1481	12.3	12 049	161	19.9	811	1133	17.5	6 459	179	11.6	1536	1136	18.4	6175	127	20.3	624
Total	2592	12.8	20 225	10 426	17.1	61 141	3114	12.2	25 602	350	19.7	1778	2390	17.4	13 775	395	11.5	3432	1748	18.3	9569	424	20.1	2110

DALYs= disability-adjusted life years; AD= attributable DALYs; PAF= population attributable fraction; OD= observed DALYs; NA= not applicable.

Note: AD and OD are numbers; PAF is a percentage.