

Table 2 Distribution of cytopenia in the study group of treatment-naïve HIV-infected patients with and without AIDS

Variable	Total (n = 493)		AIDS (n = 313)		No AIDS (n = 180)		P-value
	No.	%	No.	%	No.	%	
Whole sample							
Males	177	35.9	116	33.6	61	41.2	0.131
Females	316	64.1	229	66.4	87	58.8	0.131
Anaemia							
Males	109	61.6	74	63.8	35	57.4	0.503
Females	135	42.7	115	50.2	20	23.0	0.000
Total	244	49.5	189	54.8	55	37.2	0.000
Leukopenia							
Males	9	5.1	7	6.0	2	3.3	0.676
Females	18	5.7	14	6.1	4	4.6	0.808
Total	27	5.5	21	6.1	6	4.1	0.498
Thrombocytopenia							
Males	10	5.7	7	6.0	3	4.9	1.000
Females	12	3.8	11	4.8	1	1.2	0.244
Total	22	4.5	18	5.2	4	2.7	0.319
Anaemia + leukopenia							
Males	3	1.7	3	2.6	0	0.0	0.510
Females	4	1.3	2	0.9	2	2.3	0.699
Total	7	1.4	5	1.5	2	1.4	1.000
Anaemia + thrombocytopenia							
Males	1	0.6	1	0.9	0	0.0	1.000
Females	3	1.0	2	0.9	1	1.2	0.969
Total	4	0.8	3	0.9	1	0.7	1.000
Leukopenia + thrombocytopenia							
Total	0	0.0	0	0.0	0	0.0	–
Pancytopenia							
Total	0	0.0	0	0.0	0	0.0	–