

**Table 2 Type of foods tested and their rate of contamination with *Listeria* spp.**

Type of foods	Total samples	Contamination with <i>Listeria</i> spp.		Contamination with <i>L. monocytogenes</i>		<i>P</i> -value <sup>c</sup>
	No.	No.	%	No.	%	
<b><i>Dairy products</i></b>						
Ice cream	67	1	1.5	0	0.0	0.007
Milk	59	6	10.2	0	0.0	
Cheese	59	0	0.0	0	0.0	
<b><i>Meat products</i></b>						
<i>Kielbasa</i> <sup>a</sup>	58	4	6.9	2	3.5	< 0.001
Sausages	56	3	5.4	0	0.0	
Minced meat	73	44	60.3	1	1.4	
<b><i>Ready-to-eat foods</i></b>						
<i>Oloveyh</i> salad <sup>b</sup>	11	2	18.2	0	0.0	0.08
Fruit juice	55	1	1.8	0	0.0	
Green salad	92	5	5.4	0	0.0	
<b>Total</b>	530	66	12.5	3	0.6	< 0.001

<sup>a</sup>Type of sausage, usually served uncooked; <sup>b</sup>Mayonnaise-based salad containing cooked chicken meat, potatoes, sour cucumber and green beans.

<sup>c</sup>Chi-squared (Fisher exact) test.