

Table 4 Distribution of Iranian 4th-year dental students with experience of using tobacco products by the number of days on which the students had used tobacco products during the previous 30 days, by sex

Tobacco product/days of use in previous month	Both sexes		Females		Males	
	No.	%	No.	%	No.	%
Cigarettes: days of use						
0	75	68.2	38	86.4	37	56.1
1–5	11	10.0	3	6.8	8	12.1
6–19	9	8.2	0	0.0	9	13.6
20–30	15	13.6	3	6.8	12	18.2
Total (ever users)	110 ^a	100.0	44	100.0	66 ^a	100.0
Waterpipes: days of use						
0	112	68.7	69	78.4	43	57.3
1–5	44	27.0	19	21.6	25	33.3
6–19	7	4.3	0	0.0	7	9.3
20–30	0	0.0	0	0.0	0	0.0
Total (ever users)	163 ^{a,b}	100.0	88 ^b	100.0	75 ^a	99.9
Other tobacco products: days of use						
0	23	76.7	2	100.0	21	75.0
1–5	5	16.7	0	0.0	5	17.9
6–19	2	6.7	0	0.0	2	7.1
20–30	0	0.0	0	0.0	0	0.0
Total (ever users)	30	100.0	2	100.0	28	100.0

^aOne man did not answer this question; ^bOne woman did not answer this question.