

Table 3 Clinical location, form and number of active lesions of 137 confirmed cases of cutaneous leishmaniasis in Borujerd county, western Islamic Republic of Iran, 1998–2010

Characteristics	Males		Females		Both sexes	
	No.	%	No.	%	No.	%
Location						
Upper limbs	86	68.3	9	81.8	95	69.3
Lower limbs	57	45.2	4	36.4	61	44.5
Face	5	4.0	8	72.7	13	9.5
Neck	4	3.2	0	0.0	4	2.9
Ears	3	2.4	0	0.0	3	2.2
Upper part of chest	3	2.4	0	0.0	3	2.2
Backside	2	1.6	0	0.0	2	1.5
Abdomen	1	0.8	0	0.0	1	0.7
Shoulder	1	0.8	0	0.0	1	0.7
Clinical form of the disease						
Dry sore	74	58.7	5	45.5	79	57.7
Wet sore	52	41.3	6	54.5	58	42.3
No. of active lesions						
1	79	62.7	5	45.5	84	61.3
2	28	22.2	3	27.3	31	22.6
3	7	5.6	1	9.1	8	5.8
4	3	2.4	0	0.0	3	2.2
5	3	2.4	1	9.1	4	2.9
6+	6	4.8	1	9.1	7	5.1
All	126	100.0	11	100.0	137	100.0