

Coronavirus Disease 2019 (COVID-19) Situation Report



Weekly Report No. 270 - Saudi Arabia

23-29 December, 2021

	Global		Eastern Mediterranean Region	
	COVID-19 Cases	COVID-19 Deaths	COVID-19 Cases	COVID-19 Deaths
Current	281,808,270	5,411,759	17,122,988	315,400
Last Week	275,233,892	5,364,996	17,049,032	314,247

Saudi Arabia

	Confirmed Cases	Recovered Cases	Deaths	Active Cases	Critical Cases	PCR Tests
Total	554,665	231	8,874	4,403	43	33,049,929
in 7 days						
22/12/2021	287	113	1	2,357	35	63,737
23/12/2021	332	121	1	2,585	36	78,214
24/12/2021	325	117	1	2,792	32	64,307
25/12/2021	389	124	1	3,056	33	66,375
26/12/2021	524	142	1	3,437	40	75,682
27/12/2021	602	147	1	3,891	39	80,804
28/12/2021	744	231	1	4,403	43	84,060

Vaccination in Saudi Arabia

Total Doses Administered	Total of 1 Dose	Total of 2 Doses	Total of Booster Doses
50.5 million	25 million	23 million	2,330,999

HIGHLIGHTS

- Regions with the highest new infections over the past 7 days: Makkah followed by Riyadh.
- Saudi Arabia re-imposed wearing face masks and applying social distancing measures in all public and closed areas, including the Grand and the Prophet's Mosque starting from 7:00 AM on December 30.
- MoH: Spike in coronavirus cases is linked to Omicron. the Omicron mutant in the Kingdom is increasing.
- Dr. Al-Quwazani: 5.4% is the percentage of the mutant "Omicron" in the Kingdom.
- MoH: The booster dose of COVID-19 is available now for 16 years and above after 3 months from the second dose. Half dose of Pfizer vaccine has been approved for children and that will be administered in 2 phases.
- Ministry of Interior records 1,880 violations against precautionary measures nationally in 1 week.
- WHO launches recommendations on mask use by health workers, in light of the Omicron variant of concern: WHO interim guidelines, 22 December 2021, see link
- WHO issues an updated interim statement on COVID-19 booster doses, see link.
- WHO issues COVID-19 infection prevention and control living guideline: mask use in community settings, see link.
- WHO Updates guidelines on the management of multisystem inflammatory syndrome in children associated with COVID-19 MIS-C
- WHO publishes recommendations on the Co-administration of seasonal inactivated influenza and COVID-19 vaccines, see link.
- WHO publishes holding gatherings during the COVID-19 pandemic: WHO policy brief, see link.

IMPORTANT LINKS

- **MoH COVID-19 updates:** <https://twitter.com/saudimoh>
- **WHO's COVID-19 global situation reports:** <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>
- **WHO's COVID-19 dashboard:** <https://covid19.who.int/>
- **MoH COVID-19 dashboard:** <https://covid19.my.gov.sa/ar/Pages/default.aspx>
- **WHO recommendations on mask use by health workers, in light of the Omicron variant of concern: WHO interim guidelines, 22 December 2021:** https://www.who.int/publications/i/item/WHO-2019-nCoV-IPC_Masks-Health_Workers-Omicron_variant-2021.1
- **WHO interim statement on COVID-19 booster doses:**
[Interim statement on booster doses for COVID-19 vaccination \(who.int\)](https://www.who.int/publications/i/item/interim-statement-on-booster-doses-for-covid-19-vaccination-(who-int))
- **WHO issues COVID-19 infection prevention and control living guideline: mask use in community settings:**
https://www.who.int/publications/i/item/WHO-2019-nCoV-IPC_masks-2021.1
- **WHO updates guidelines on the management of multisystem inflammatory syndrome in children associated with COVID-19**
[WHO issues guidelines on the treatment of children with multisystem inflammatory syndrome associated with COVID-19](https://www.who.int/publications/i/item/WHO-2019-nCoV-IPC_masks-2021.1)
- **WHO recommendations on the co-administration of COVID-19 and influenza vaccines:** [Coadministration of seasonal inactivated influenza and COVID-19 vaccines \(who.int\)](https://www.who.int/publications/i/item/co-administration-of-seasonal-inactivated-influenza-and-covid-19-vaccines-(who-int))
- **Holding gatherings during the COVID-19 pandemic: WHO policy brief:**
<https://www.who.int/publications/i/item/holding-gatherings-during-the-covid-19-pandemic-who-policy-brief-2-august-2021>

IMPORTANT DEVELOPMENTS

The World Health Organization issued an emergency use listing (EUL) for Nuvaxovid™





The new vaccine was developed by Novavax and the [Coalition for Epidemic Preparedness Innovations](#) (CEPI), and is the originator product for the Covovax™ vaccine that received WHO emergency use listing on 17 December.

Both vaccines are made using the same technologies. They require two doses and are stable at 2 to 8 °C refrigerated temperatures.

The Novavax vaccine (NVX-CoV2373) consists of a recombinant SARS-CoV-2 spike protein nanoparticle administered as a co-formulation with the adjuvant Matrix-M. Protein-based vaccines have been used against diseases such as pertussis, human papillomavirus, and hepatitis B. Matrix-M is a novel adjuvant that has been used in studies but has not previously been used in any licensed vaccine.

The efficacy of NVX-CoV2373 has been assessed in three phase 2 and phase 3 trials involving participants aged 18 years or older. In a phase 3 study conducted in the United Kingdom during a period in which the SARS-CoV-2 Alpha variant was predominant, vaccine efficacy (VE) against mild, moderate, or severe COVID-19 was 90% (95% CI: 80–95) from 7 days after the second vaccine dose, with a median follow-up of 56 days after the second dose. VE against mild, moderate, or severe disease in persons less than 65 years of age was 90% (95% CI: 80–95) and in those 65 years and older 89% (95% CI: 20–100). Studies of NVX-CoV2373 have demonstrated an acceptable safety and reactogenicity profile in adults ≥18 years of age, detailed data on the efficacy and safety of this vaccine can be found in the background document on the NVX-CoV2373 vaccine (see WHO website). The data reviewed by WHO support the conclusion that the known benefits of NVX-CoV2373 outweigh the risks that are known or considered possible. Therefore, WHO recommends the use of NVX-CoV2373 in persons aged ≥18 years. As sufficient vaccine supply will not be immediately available to immunize all who could benefit from it, countries are recommended to use the WHO Prioritization Roadmap and the WHO Values Framework as guidance for prioritized vaccine use, based on population subgroup.

The recommended primary vaccine series is two doses given intramuscularly into the deltoid muscle at an interval of 3–4 weeks. The vaccine should not be administered with an interval of less than 3 weeks. WHO is currently assessing the need for and timing of booster doses. Data on the duration of continued protection are currently still missing.



IMPORTANT CONTACTS

- The National Focal Person for COVID-19 is Dr Abdullah Asiri, Assistant Deputy for Preventive Health, MoH, email: AbdullahM.Asiri@moh.gov.sa